



Primary Care in LMIC Cities 2

Policy and service delivery proposals to improve primary care services in low-income and middle-income country cities

Richard J Lilford*, Benjamin Daniels*, Barbara McPake, Zulfiqar A Bhutta, Robert Mash, Frances Griffiths, Akinyinka Omigbodun, Elzo Pereira Pinto Jr, Radhika Jain, Gershim Asiki, Eika Webb, Katie Scandrett, Peter J Chilton, Jo Sartori, Yen-Fu Chen, Peter Waiswa, Alex Ezech, Catherine Kyobutungi, Gabriel M Leung, Cristiani Machado, Kabir Sheikh, Sam I Watson, Jishnu Das



The landscape of primary care services in low-income and middle-income country cities is diverse and dynamic, yet the quality of care received is too often low and the financial cost to the patient high. In the second Paper in this Series, we argue that shaping the primary care market is likely to provide larger returns to scale than individual quality improvement initiatives. Among other things, the market can be shaped by regulation and targeted public investment to crowd out poor providers and motivate those that remain to improve. Additional supply-side initiatives for which there is evidence include measures to educate and motivate the workforce, skill substitution and formation of clinical primary care teams, information technology, and improving the supply of medicines and diagnostics. Demand-side measures include reducing out-of-pocket expenses and promoting health literacy and user advocacy. Research is urgently needed into access for people who are unregistered (eg, those who sleep on the streets), those in peri-urban areas and towns, and on cost-effectiveness, and sustainability of beneficial interventions.

Introduction

This is the second and final paper in a Series about the provision of primary care services in urban settings in low-income and middle-income countries (LMICs). In the first paper, we described the existing landscape of access, use, cost, quality, and choice in primary care. The evidence summarised there described a dense, diverse, and dynamic landscape of primary care provision in LMIC cities, where most community clinics have significant excess capacity. The average quality of care was low, and out-of-pocket expenditures could be substantial, but there was considerable variation in both. Thus, the supply and demand for primary care in these environments resembled markets for commoditised services, rather than top-down publicly mandated health systems. Here, we build on that evidence to discuss what might be done to improve primary care provision in LMIC cities and explore the potential for success among several commonly proposed strategies, recognising that the existence of this market structure provides a set of options to the urban policy maker that are not available in a typical rural setting.

In this Series, we reviewed evidence regarding policy or service interventions to improve primary-care quality in LMICs and situated these within a conceptual framework that reflects on how each intervention interacts with the broader structures of competition and choice in the urban environment. As in the first paper in this Series, we use the Institute of Medicine classification of outputs to which a health system should aspire (effectiveness, safety, patient-centredness, accessibility, efficiency, and equity).¹ We use system performance to refer generically to improvement across the Institute of Medicine criteria, and quality when we want to refer to clinical care consultations requiring the clinician to recognise conditions, recommend appropriate actions, and provide

patient-centred care. The method of literature retrieval used to inform our analysis is described in panel 1.

Conceptual framework to classify the interventions to improve system performance

There are numerous frameworks describing the determinants of system performance, including the WHO–European Observatory document on health system

Key messages

- Although services can be improved at the supply and demand sides, the existence of a market in urban areas opens another route to improved services not typically available in rural areas—namely, the possibility to shape the market.
- At the supply side, there are compelling reasons to develop primary care teams incorporating community health workers.
- Other promising supply side interventions include education, management training, and shared records (starting with patient-held paper records as precursors to electronic records).
- Demand-side interventions to improve self-confidence and health literacy are effective interventions that align well with market shaping.
- Policy makers have additional powerful strategies to shape the market in the form of strategic investment in public services and regulation. These strategies have been shown to both crowd out poor-quality providers and crowd in better quality providers.
- Research on primary care in low-income and middle-income country cities is at an early stage and should more often consider the opportunity cost of interventions than at present.

Lancet Glob Health 2025;
13: e954–66

See [Comment](#) page e795

*Joint first authors

This is the second in a [Series](#) of two papers about primary care in LMIC cities

Institute of Applied Health Research, University of Birmingham, Edgbaston, UK (Prof R J Lilford PhD, K Scandrett MSc, P J Chilton BSc, J Sartori BA, S I Watson PhD); Department of Global Health and Population, Harvard T H Chan School of Public Health, Boston, MA, USA (B Daniels MSc); The University of Melbourne, Parkville, VIC, Australia (Prof B McPake PhD); Institute for Global Health & Development, The Aga Khan University, South-Central Asia, East Africa, and UK, Karachi, Pakistan (Z A Bhutta PhD); Centre for Global Child Health, The Hospital for Sick Children, Toronto, ON, Canada (Z A Bhutta); Department of Family & Emergency Medicine, University of Stellenbosch, Cape Town, South Africa (Prof R Mash PhD); Warwick Medical School, University of Warwick, Coventry, UK (Prof F Griffiths PhD, Y-F Chen PhD); Centre for Health Policy, School of Public Health, University of the Witwatersrand, Johannesburg, South Africa (Prof F Griffiths); College of Medicine, University of Ibadan, Ibadan, Nigeria (A Omigbodun MD); Center of Data and Knowledge Integration for Health, Oswaldo Cruz Foundation–Brazil, Salvador, Brazil (E P Pinto Jr PhD); Global Business School for Health, University College London, London, UK (R Jain ScD); African Population and Health Research Center, Nairobi, Kenya (G Asiki PhD, C Kyobutungi PhD); University Hospitals

Birmingham NHS Foundation Trust, Birmingham UK (E Webb MBBS); School of Public Health, College of Health Sciences, Makerere University, Kampala, Uganda (Prof P Waiswa PhD); Dornsife School of Public Health, Drexel University, Philadelphia, PA, USA (Prof A Ezech PhD, Prof K Sheikh PhD); School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region, China (Prof G M Leung MD); Sergio Arouca National School of Public Health, Oswaldo Cruz Foundation, Rio de Janeiro, Brazil (Prof C Machado PhD); McCourt School of Public Policy and the Walsh School of Foreign Service, Georgetown University, Washington, DC, USA (Prof J Das PhD)

Correspondence to: Prof Richard J Lilford, Institute of Applied Health Research, University of Birmingham, Edgbaston B15 2TT, UK r.j.lilford@bham.ac.uk

See Online for appendix

Panel 1: Search strategy and selection criteria

We reviewed papers retrieved for the first paper in this Series that included evidence regarding policy or service interventions (henceforth, policy interventions), including Cochrane overviews²⁻⁵ and other systematic reviews dealing specifically with quality improvement in low-income and middle-income countries, including the review by Rowe and colleagues,⁶ which collated and summarised outcomes from health system interventions before 2016 from 337 studies. We include scoping and systematic reviews that evaluate policy interventions, including those covering specific countries,⁷ specific regions (Asia-Pacific),^{8,9} children,¹⁰ and specific policies such as payment for performance,¹¹ mobile health,¹² paper records,¹³ delivery of medicines,¹⁴ and cost effectiveness studies.¹⁵

A considerable body of evidence on policies to improve health care was recently compiled by Rowe and colleagues.⁶ This evidence covers urban and rural settings. We, therefore, performed a re-analysis of this existing systematic review to obtain data in a specific urban context. We obtained an updated version of the underlying database, which we filtered to include only studies conducted in urban, peri-urban, and mixed urban-rural settings, either in a primary care community health facility, or hospital general outpatient departments (which we termed polyclinic). This filtering yielded a list of 74 studies (appendix p 8), which we checked to confirm appropriateness to this paper's topic.

performance,¹⁶ and the study by Lilford and colleagues cited in the first paper of this Series,¹⁷ but we use a standard classification of factors that turn inputs into services. That classification, represented in the figure, covers a number of different areas: measures to improve the output of an existing set of clinical facilities servicing an existing set of care-seekers, without changing the composition of provider facilities—these could be called supply side interventions; consumer empowerment and pressure at the level of the community or individual patients, including both financial and informational elements that redirect consumer decisions and willingness to pay—these could be called demand-side interventions; and measures to change the composition of clinical facilities using policy levers, including, but not limited, to investing in public services and regulation—these could be called market-shaping interventions.

Supply side measures to improve the performance of an existing set of providers

One approach to improving primary care providers' performance, widely represented in the literature, is to intervene to enhance care quality delivered by existing providers (figure). Conceptually, the performance of a given set of service providers can be affected by three broad categories of intervention: firstly, by improving performance directly by increasing knowledge

or skills (eg, through education and training) or by changing attitudes or motivation (eg, through education and incentives); secondly, by increasing the availability of complementary resources for those providers, through improvements to facilities, amenities, or commodities such as medicines and diagnostics; or finally, by allocating tasks more efficiently among teams of providers who have various comparative advantages.

Improving the performance of individual providers

Educational programmes to increase medical competence

Medical knowledge deficits among practitioners contribute to poor quality care, but are not the main or only reasons according to the ten-country sub-Saharan African study described in the first paper in this Series.²² There is extensive literature on educational interventions to improve primary care in LMICs, which generally shows that they are effective, but that the magnitude of effect is modest. 40 of the 74 LMIC city studies retrieved in our literature search (appendix p 8) included an educational intervention (in 15 studies this was the only intervention, while in 25 it was one of several interventions). In panel 2, we provide a set of evidence-based principles that should be considered when designing educational programmes.⁶

Financial incentives for improved provider performance

The well documented know-do gap in quality measurement literature suggests that even skilled providers do not put a large share of their medical knowledge into practice. Some of this underperformance can be put down to poor motivation or effort. Poor motivation is an archetypal indication for incentivisation. One commonly suggested method of incentivising providers to do what they know is to make a payment at clinical level for improved performance. However, others have theorised or found that financial incentives can displace intrinsic motivation and replace valuable activities not included in the incentive.³³⁻³⁵

Systematic reviews do not provide unequivocal evidence in favour of payment for performance. A review in high-income countries³⁶ found that payment for performance provided inconsistent results and observed that the incentivised action displaced other, worthwhile, actions. A Cochrane systematic review of payment for performance specifically in LMICs found that "the effects...on delivery and use of services is mixed overall"³⁷ and most studies were judged to be of uncertain quality.^{11,37} In the Cochrane review, only 17 of the 59 reviewed studies compared payment for performance with a comparator investment, while 42 compared the payment for performance intervention with service as usual rather than with alternative uses of a similar amount of funding. In contrast, a World Bank study compared payment for performance with direct facility financing (DFF) in five sub-Saharan African countries. Again, payment for performance "adds little gain over




Figure: Classification of interventions to improve primary care services

Note, we do not discuss macro policies,^{38–20} such as training the medical and nurse workforce, fiscal policy, or the built environment, that affect the whole health-care system, rather than primary care services in particular. Much of this material is covered in WHO reports²⁰ and a previous Commission in *The Lancet Global Health* on financing primary health care.²¹ At the opposite end of the chain, we do not discuss micro policies (interventions) targeted at specific clinical areas such as tuberculosis, vaccination, and mental health.

DFF”.³⁸ That said, a widely cited study (included in the Cochrane review) that compared payment for performance with provision of a similar amount of money for general use, observed positive effects of the intervention (on some outcomes),³⁹ and a cluster trial from Brazil published after the review found that payment for performance improved care, especially in impoverished areas.⁴⁰ Overall, we are circumspect about the value of payment for performance,⁴¹ and note that there is a practical limit to the number of such initiatives to which clinicians can respond.

Supplies and facilities essential to good practice

Medical supplies (medicines, vaccines, diagnostics, equipment, and aids)

We observed—as noted in the first paper in this Series—that, without a large generics industry, as in Bangladesh,⁴² some of the biggest problems for low-resource city dwellers are the cost, unavailability, and quality of medicines.^{42,43} However, this problem is tricky because the supply chain is complex and a market clearing price should be set so that supply is ensured while keeping

costs low to both patients and the public purse. Numerous approaches have been tried,⁴⁴ including public subsidy, where, as countries become richer, they can increasingly adopt financing mechanisms that cover costs more comprehensively, and pooling savings by implementing policies where service users club together to increase their purchasing power.⁴⁵

Regulation is another approach; pharmacies are regulated in most LMICs, but regulation is poorly enforced, reducing its use as a policy instrument, for example to ensure suppliers hold minimum stock levels.⁴⁶ Simultaneously, there are arguments to regulate against allowing clinics to both diagnose and prescribe to avoid the situation whereby providers have a perverse incentive to overprescribe the most expensive medicines (and tests). Such regulation is possible in cities where there are plentiful pharmacies and laboratories, but not in rural areas where access would be restricted.

Finally, there is the issue of supply chain management combined with poor overall stock control and supply chain methods, and poorly written contracts, which lead to poor maintenance of equipment and stockouts of

Panel 2: Measures to capitalise on continuing professional development opportunities

- The education programme should be carefully designed, starting with a needs assessment.²³
- Training topics should include provision of person-centred care and use of appropriate referral pathways to reduce documented delay in diagnosis of diseases such as cancer.²⁴
- Front-line interventions are most effective when backed up by supportive changes higher in the management hierarchy.⁶ Education is an archetypal example as educational interventions can be sustained and knowledge reinforced²⁵ if they are incorporated in the continuing professional development frameworks that now exist in most low-income and middle-income countries.^{18,26}
- As much as possible, learning should be active rather than didactic, and should promote problem-solving and self-directed learning.²⁷ A 2023 non-randomised study in China showed that small group learning and use of standardised patients to provide constructive feedback improves performance and patient satisfaction with subsequent consultations compared with more passive teaching methods.²⁸
- Instructors should provide motivation and inspiration to help close the knowledge–practice gap and encourage team working and collaboration.^{29–31}
- Those who commission education should be alert to the risk that educational providers will exploit any monopoly power to charge high fees, a problem described in the Philippines.²⁶
- Programmes should be developed iteratively and evaluated wherever possible.^{23,32}

vaccines and medicines.⁴⁷ A 2021 scoping review evaluated centralised dispensing and packaging systems for people with chronic diseases. The systems have used a range of alternative delivery mechanisms (ranging from smart lockers to home delivery), and have improved affordability, adherence, and reduced demand on facilities.¹⁴

Information systems and digital integration

There are a number of ways that information systems and digital integration could be used to improve practice, starting with monitoring quality. Routinely collected data can be used to help an organisation monitor its performance, covering criteria such as vaccination rates, antenatal clinic attendances, and follow-up of patients with HIV or hypertension. Such data can be used to audit performance against indicators, hence acting as a stimulus for improvement. The problem is developing a system for measurement of the necessary performance criteria, given that data collection systems are currently rudimentary throughout

most LMICs, with exceptions such as Brazil.⁴⁸ The private sector in some countries collects data to enable billing and insurance claims, but not to purposefully measure the quality of care.⁴⁹ The development and implementation of universal health coverage could act as a stimulus to implement data collection. In time, such systems could incorporate patient electronic records.

Clinical records could be used to support clinical care. Modern health care requires that an individual is served by many providers, and in many places there is no system for information sharing.⁵⁰ Since relational continuity is inevitably interrupted, informational continuity is essential,^{51,52} especially in cities where care is shared across a large number of providers. Although a system in Western Cape, South Africa, based on an inference or probabilistic approach to data linkage has achieved success in integrating fragmented individual electronic health data,⁵³ this is an exception and the difficulties in digitising notes are consistently underestimated.^{54,55} There is a tendency to think immediately of electronic patient records,^{21,56,57} but structured patient-held paper records⁵⁰ are an effective, but underused,⁵⁸ means of information capture and transfer according to a 2022 systematic review.⁵⁹ Pending development of effective digital solutions, we strongly and urgently advocate use of structured patient-held notes to improve informational continuity, as in the case of Kenya (and worldwide in the specific case of maternity care).

Virtual (remote) consultations (not to be confused with chatbots) are increasingly common worldwide and were vital during the COVID-19 pandemic. Our 2021 umbrella review of seven systematic reviews of this intervention in (mostly urban) LMICs all show patients benefiting.^{55,60} There is, however, massive headroom to expand use of this innovation in city areas where people are employed in the informal sector and lose income when attending a clinic. Previous research has produced guidance on how mobile consulting or chatbots can be structured and deployed so people who are low-income and those who are marginalised are not excluded.^{61,62}

There is an extensive literature on use of apps and messaging systems to provide health messaging to people—for example, medication reminders or diabetes self-management.⁶³ Mobile phones are highly available in LMIC cities, even in slum areas. However, only 88 of the 5322 households in a study of seven slums across four countries,⁴² reported using their phone to access and receive health information, advice, or care in the previous 12 months.⁶⁰

Improving organisations to enable more efficient service management

Primary care teams

In the first paper of this Series, we highlighted the crucial role that primary care services play in connecting communities with specialist care.⁴ We also mentioned

(with examples from Brazil and South Africa) that these teams frequently include community health workers (or equivalents). Achieving national coverage with a network of such teams takes time—as shown by Kazakhstan’s two decade process.⁶⁴ However, there are good arguments in favour of policies to gradually substitute multiple, poorly integrated providers for primary care teams, summarised in panel 3.

Vertical systems of care for specific conditions, such as tuberculosis, HIV, and maternal care, have been successful, but they cannot be provided for all conditions separately, and have limitations in terms of co-ordination for continuity across multiple chronic diseases.⁷² The 2024 WHO primary health care global report is clear that some countries (such as Brazil, where the public sector plays a dominant role) have been more successful than others (such as India, where it does not) in transferring resources to primary care clinics and teams.⁷² However, there are policy options for implementation of such teams—particularly whether they are incorporated in a hospital system and, if not, whether they act as gatekeepers in a system where access to hospital care is controlled. A study from China that compared these options favoured primary care teams that were financially and managerially independent of local hospitals.⁷³ Lastly, we cannot assume that the quality of care is automatically good in primary care teams according to evidence from South Africa⁷⁴ and China.⁷⁵

Management support and supervision

One of the advantages of clinical teams over very small or single-handed clinics is that ready-made teams are available to receive the training and managerial support. Managers can be trained and coached to provide an environment supportive of improved performance and produce systemic change in attitudes. The effects of such management support in primary care in LMICs have been reported in recent studies,^{76–79} including an 81 study systematic review of supervision.⁸⁰ These articles show generally positive effects, especially when supervision involves group problem solving. It is possible that regulation is most effective and sustainable when linked to more collaborative and supportive management styles.^{81–84} The point we make here is that management training is a kind of education trying to change motivations and attitudes. Also, attitudes and motivation are key to clinical care—for example, the knowledge–practice gap described above. The corollary is that management or leadership training and clinical training are connected—they should be integrated,^{85–87} as in the approach to improve primary cancer care in sub-Saharan Africa.⁸⁸

Facilitating demand through consumer empowerment

It is important to implement measures to enable or encourage people to choose higher-quality providers

Panel 3: Advantages of creating a system of primary care teams based in primary care facilities and supporting community health workers

- Primary care teams, based in health centres or clinics, are a vehicle to integrate facility and community care. Incorporating community health workers within the primary care team offers an effective strategy for disease prevention^{65,66}—for example, in ensuring more equitable vaccine coverage⁶⁷ and providing a direct route to medical care if a community health worker comes across a malnourished child or adult with chronic cough that might be caused by tuberculosis or cancer. However, it is important that roles and responsibilities are delineated explicitly.⁶⁸
- Teams can provide the necessary coordination that is often missing in a fragmented system described in the first paper in this Series. Community health workers are often embedded in the community, have greater access to marginalised groups, and are socially acceptable. Their presence can also facilitate early discharge to the community, management of chronic disease and mental health, rehabilitation, and palliative care in line with WHO Disease Commodity Packages principles.⁶⁹ Thus, whereas a survey of returning patients in private clinics in Nairobi, Kenya, showed that care for such conditions is poorly co-ordinated,⁷⁰ a study of primary care teams in South Africa found much higher scores for co-ordination, continuity, and comprehensiveness.⁷¹
- Staff absences can be covered, thereby improving continuity of care.
- Teams form the framework around which clinical education and managerial support can be delivered.
- Teams form the vehicle for skill substitution (for example, where members of the team effectively take over tasks often associated with doctors). A 2018 systematic review showed that such substitutions are cost effective.¹⁵
- Teams provide an environment where the doctor is available when needed.⁷⁹

(figure). In essence, the market described in the first paper of this Series should be augmented to function ever more efficiently by empowering people financially, better informing them, and facilitating collective action and advocacy in shaping services.

Improving the affordability of primary care services

Remove or reduce user fees for public services

As shown in the first paper in this Series, user fees tend to suppress demand,⁸⁹ including for preventive care and particularly for children. A systematic review of financing interventions in the Asia–Pacific region (covering two-thirds of the world population) found that removing user fees improved care when public facilities were remote or of very poor quality.⁸ A randomised trial of removing user fees for children in Ghana not only resulted in increased

use of public services but also improved health outcomes,⁹⁰ whereas a study in Malawi showed that user fees had a particularly disadvantageous effect on children.⁹¹ Thus, there is a strong argument for reducing user fees by governmental and non-governmental organisations whenever possible.⁹²

Contracting private providers from public funds

There are many examples where copayments are provided for independent providers to provide services, either directly or by providing vouchers or insurance for patients. These have been successful in reducing user fees and increasing equality of access and quality in Uganda,⁹² improving equity and care of primary care in the Asia-Pacific region,⁸ improving diabetes care in Mexico,⁹³ and reproductive care, again in the Asia-Pacific region.⁹⁴ Likewise, four social insurance schemes examined by the World Bank improved access to various degrees, depending on how generous they were.⁸ Additionally, there are examples where the private and public sectors have collaborated and produced promising results, such as contracting out primary care service to non-governmental organisations in the Urban Primary Health Care Project by the Government of Bangladesh,⁹⁵ working with private doctors with managerial and technological support from a non-governmental organisation in the Mohalla Clinics of Delhi,⁹⁶ and implementing agreed referral pathways in Kenya.^{97,98}

Based on economic theory, providers in urban environments might be more responsive than those in rural environments to the incentives provided by copayments since they work in a competitive environment. Therefore, the purpose of such contracting might not be to generally increase the availability of services, but to harness scarce information about quality to drive demand by strategically subsidising or implicitly endorsing better-performing providers.

Community financing

Community financing is a term coined in the 1980s⁹⁹ to include self-help forms of financing strategies including revolving drug funds, community health insurance, and savings groups. These mechanisms pool local resources¹⁰⁰ to support the costs of health-care for those who fall sick.

Initial enthusiasm for initiatives such as the UN Bamako Initiative¹⁰⁰ waned in the 1990s after funds contributed were found to be insufficient to sustain the regular supply of medicines.¹⁰¹ More positive evidence associated with newer models has emerged more recently in the form of large prescription savings groups,⁴⁵ since adopted by Nairobi County, Kenya. In these models, the collective savings for the group used to procure medicines at a wholesale price make medicines more affordable to the patients. Nevertheless, this model is a novelty and small-scale user fees,

alongside limited voluntary insurance models, are the more common interpretation of the term community financing globally, and are more commonly features of rural than urban areas.^{102–104} These models tend to result in the exclusion of the poorest, unless subsidised, and sometimes even if subsidised.^{101,105–107}

Lastly, there has been interest in savings groups as a poverty alleviation method in development economics. Of particular interest here are initiatives that combine community action groups to provide peer support and improve health for chronic conditions with micro-economic initiatives. These have been used in rural areas to support people affected by leprosy,¹⁰⁸ and, more recently, a pilot study of a self-help group for patients with hypertension, diabetes, or both has produced promising results in a slum in Nairobi, Kenya. (Asiki G, et al., unpublished)

We do not have space here to consider the large and complex literature on conditional cash transfers that incentivise specific behaviours such as attending antenatal clinics or vaccination, nor non-conditional transfers, which belong in the general realm of development economics.^{109,110}

Informing and empowering patients to make better choices

Community and peer self-support groups might be of value even if they do not include financing activities as a feature of self-help groups. Such peer support or self-care groups provide psychological support and share knowledge to improve health literacy and help people navigate the health-care system, guiding people to registered (ie, qualified) providers, well supplied medicine outlets, and appropriate specialist services. Such groups can build on existing support networks that exist, for example, in slum communities.¹¹¹ There is a large amount of literature on such groups that is mainly positive, but a systematic review of seven cluster trials¹¹² provides some evidence that they work better in rural areas than cities—perhaps because people in cities more often have to leave home to work in the informal sector, and therefore cannot easily participate in community groups.⁹⁷ Likewise, there is literature on the use of paid navigators to support people, especially those with low health literacy, to overcome barriers on the health-care pathway.¹¹³

Moving beyond informing and supporting individuals to access appropriate health care, communities can help shape services that affect them in line with WHO people-centred and integrated care principles.¹¹⁴ Following this principle will help ensure that services meet diverse needs provided that the most vulnerable, including unregistered migrants, are included.

Improving the composition of the provider population (market shaping)

As noted in the first Paper in this Series, the profusion of small, mostly private, clinics is associated with poor

quality services. This is evidence of market failure and could even lead to calls for an outright ban on private care. However, we reviewed the evidence (observational and experimental) in the first paper in this Series and found no clear winner in the public versus private debate. As argued in a previous Series on universal health coverage,¹¹⁵ there is no good argument to ban private practice outright. The problem then is to improve quality without reducing access.

In this section we discuss policies to manage or shape the market, rather than abolish it, by exerting influence over the quality of the provider population (figure). Specifically, this type of approach is predicated on the idea that policy can either encourage new higher-quality providers to enter the market or cause existing lower-quality providers to exit, while maintaining a large enough provider population that access, choice, and competition are not compromised. Two (non-exclusive) options present themselves to shape the market.

Investing in public services

In low-density settings, natural monopolies are likely to emerge in primary care provision, and therefore direct public service provision (much like a utility) is often viable as the main policy approach. In dense environments, however, the market will offer diversity, and the expansion of public services should be used strategically to influence the choices of private provider facilities, as well as of care-seekers, with knock-on effects for outcomes from the mixed public and private market overall.

There are numerous arguments that public investment is not only a benefit in and of itself, but that it also has market-shaping potential. First, this idea is in line with the economic theory that high-quality services crowd out poor-quality providers while encouraging those that remain to improve. Second, there is evidence that competitive pressure can drive up quality, including in health care.^{116,117} Third, countries that spend more on public services with reasonably remunerated practitioners have fewer small-scale and informal providers.¹¹⁸ Fourth, we know that in LMIC cities the fundamental ingredient for market pressure is present since patients actively seek out high-quality providers, bypassing those of lower quality, as shown in the first paper in this Series. Finally, modest increases in Indian State health spending were associated with substantial substitutions of public for private care (especially for disadvantaged groups).¹¹⁹

In dense urban environments, there is emerging evidence that public investment does not only crowd out low-quality providers, but also that the agglomeration effects might, on net, crowd in new high-quality private services. This effect was seen with respect to hospital provision in a 40-year natural experiment in Malaysia, where political constituencies tended to acquire new public hospitals when their member of parliament was promoted to the cabinet, and where a new public hospital caused private hospitals to be built nearby.¹²⁰ The theory

is that the new public hospitals created a favourable labour market that more than compensated for the more competitive customer market. Evidence regarding regulation (panel 4) suggests that this phenomenon might also occur in primary care. In conclusion, not only is public investment in facilities likely to shape markets, but also to be more scalable and sustainable than outreach quality improvement initiatives that should be delivered to each facility repeatedly.

Regulation of private health-care provider organisations

Regulation of services with a view to improving provider quality involves a legal framework, a regulatory body, registration of provider units, setting standards, conducting inspections (audits), and enforcement. As with the crowding out theory, regulation might cause some clinics to close and others to improve.⁷⁶ Regulation could improve quality of care by a number of theoretical mechanisms: removing the worst clinics; acting as an incentive for clinics to improve; creating an environment in which new providers with the potential to provide high-quality services believe they can thrive and therefore enter the market; or providing information about where improvement is most urgently needed. Further to this last point, regulation could be coupled to improvement initiatives, as seen in the examples from South Africa and Tanzania.^{76,79}

Most countries now register and regulate formal (allopathic) providers and pharmacies. For example, in Africa, the organisation SafeCare provides regulation services across a large number of countries.¹²⁶ As with payment for performance mentioned above, regulation faces a measurement challenge: regulators should define, ascertain, and then act on measures of quality. This opens extensive possibilities for misclassifying providers (in both directions) and, as shown in the first paper in this Series, the measurement of care quality itself is a complex task. It becomes more so when it has consequences for providers who will frequently game the system by manipulating the measurements.¹²⁷ Until recently there was scant high-quality evidence on the effectiveness of regulation of primary care providers, but now two randomised controlled trials have become available, both with positive results. The literature is summarised in panel 4.

Finally, regulation and enforcement can be a powerful policy tool driving up quality and shaping the provider market, but more work is needed to find the most cost effective approach for different types of providers. Furthermore, context is important and trial evidence shows what can work, but not what will—for example, regulation can open the door to corruption. This consequence is a risk when the government has limited capacity and in cities where there are overlapping jurisdictions with different interests. We recommend that regulation is phased and evaluated and well regulated itself.

Panel 4: Evidence of effect of regulation on health markets

- In 2011, Flodgren and colleagues carried out a Cochrane systematic review into evidence regarding the effects of regulation,¹²¹ which was updated in 2016,¹²² and then by WHO as part of their recent guidance. Two evaluations of regulation were found, only one of which was in a low-income and middle-income country (LMIC)—a cluster-randomised, controlled trial of the Council for Health Services Accreditation scheme for South Africa. The intervention was based on an inspection visit, including assisting in enabling the facility to improve on sub-optimal standards. This study showed a marked improvement in practice in the intervention group compared with the control but does not provide evidence on regulation alone.
- Bedoya and colleagues reported results from their randomised trial of a regulatory reform in the health sector.¹¹⁷ The authors evaluated the 1-year effect of this intervention. To evaluate this reform, the authors divided all 1348 health facilities in three Kenyan counties into 273 markets, and then randomly allocated the markets to treatment and control groups. Government inspectors visited intervention health facilities and, depending on the results of their inspection, recommended closure, or a timeline for improvements; 25% of all private facilities were shut down by the government in the intervention group. The authors first show that the intervention improved quality of care as measured by adherence to a checklist of patient safety and structural measures, without any decline in patient loads or increase in prices. The authors found that 87% of the increase in quality was due to improvements in existing facilities, only 5% was due to an increase in closures, and the remaining 8% was due to the entry of new facilities (consistent with the crowding in hypothesis). The authors show that the latter is likely due to removal of cheaper low-quality provider facilities in a market where higher quality is rewarded through the regulatory system. The authors concluded that, in this case, instead of reducing competition in the market (as is often the case), regulation seems to have increased it.
- We found two additional randomised controlled trials involving pharmacy regulation in LMICs,^{123,124} both showing improvement in pharmacy practice, but, in each case, regulation was combined with other interventions, such as training pharmacists. A recent systematic review examined the role of regulation enforcement and various forms of education that were mostly effective in reducing over-the-counter sale of antibiotics, especially where multiple strategies were concerned.¹²⁵

Discussion

The key theme of this Series is that cities provide an environment in which it is possible to move beyond traditional clinic-level interventions and instead improve the average quality of services in urban markets by (indirectly) exerting influence over provider choices of whether and how to participate in the market. We refer to these strategies as market shaping as they primarily operate through changing the composition of the provider population, and careful stewardship of the rules of exchange. For example, in the US health market, insurers and government payers shape the market by including or excluding providers from their coverage networks, and by setting restrictions on pricing and agglomeration that affect provider decisions to offer various services within those networks.

There are several ways to shape the market. Regulation can simultaneously remove low-quality providers, improve the quality of remaining providers, and encourage new, high-quality entrants (including direct public provision). We have provided numerous strands

of evidence to support the theory that greater public investment is associated with crowding out of poor-quality providers. Evidence shows that providing high-quality services can crowd in new providers. The theory here is that a high density of providers in cities, with lower densities in towns and peri-urban areas (described in the first Paper in this Series), could be the result of crowding in through agglomeration of clinical services. The phenomenon of crowding in or agglomeration is well supported in general economics.^{128,129}

Market shaping is arguably more scalable and sustainable than quality improvement initiatives requiring direct public expenditure. Reshaping the primary care services market in cities thus has greater potential effect compared with even the most successful individual clinic-level programmes.¹³⁰

In addition to the conventional improvement strategies that improve existing services at supply and demand sides (figure), there is evidence supporting education (incorporated with continuing professional development), improving information systems (using paper-based, patient-held notes where digital technology cannot be implemented in the short term), and providing remote consultation options. Managerial support is effective, but is only applicable where teams already exist and might be difficult to scale and sustain. Conversely, strengthening primary care teams is an effective method to improve clinical practice, while providing co-ordination of prevention services and chronic care, along with its market-shaping potential. Likewise, regulation (perhaps especially if combined with management support) has both direct potential benefits alongside market-shaping potential. However, there are some interventions where we think the evidence is more equivocal, pending additional research, including payment for performance, and more research is needed regarding innovations to improve availability of equipment and medicines.

There are good reasons to inform patients to make them discriminating citizens and to mentor self-help or peer support groups. These measures contribute to market shaping. Removing user fees and providing free or discounted medicines for people with long-term conditions are priorities in overcoming important barriers to health care. However, policy makers can only remove fees for services they provide. To influence other parts of the service, services can be contracted out or public subsidies to support wider access to private services can be provided. We find these subsidies can be effective in improving access, and that revolving drug funds and community health insurance savings groups are not sustainable (or equitable) without external support. This reliance on subsidies leaves an open question for policy makers who can invest in either public services (expanding services and reducing user fees), providing subsidies, or both. It might be the case that different balances between these options are more

suitable than others according to the state of current development of services and the broader economy. The question of the role of the state in buying versus supplying services is not resolved in economies of any scale. However, in LMICs where health services are scarce or unequally distributed, expanding public provision and promoting health equity should be priorities.

It is important to recognise that policy interventions play out differently in different contexts. First, clumsy implementation where policy makers and practitioner objectives are not aligned can stymie interventions—a 2023 special series by Schneider and colleagues draws attention to potential implementation pitfalls.¹³¹ Second, interventions such as regulation and education based on continuing professional development can be misused by unscrupulous service providers, as described above. Third, interventions can induce unintended effects and back-fire; for example, payment for performance can cause priorities to be distorted, revolving drug funds can widen disparities, and removing fees for drugs can lead to prescriptions of unnecessary medicines. Fourth, an effective intervention is not necessarily cost effective; the opportunity cost should always be considered. Fifth, there are strong arguments to compare different designs within a class of interventions. For example, we have advocated primary care teams, but these can be designed in different ways—as an extension of the hospital service or independent, in which case they might or might not have a gatekeeping function. We found only one study, conducted in China, that has compared these three alternatives head-to-head.⁷³

Many studies combine interventions into compound packages—for example the BRAC Manoshi programme of community engagement,¹³² and a package involving self-care, community health worker outreach, and community engagement that was effective across a number of health outcomes in a randomised controlled trial in slums in Mumbai, India.¹³³ A combination of community activation and provision of accessible mobile outreach services resulted in improved uptake of vaccination in a large, non-randomised cluster study in Pakistan with substantial effect on reducing inequities.¹³⁴ Most urban primary care system interventions are combined, posing challenges for evaluation.

The key point is that provider and consumer incentives are jointly determined and interventions on one side can bolster the other. For example, public funding for care at private facilities, whether through sub-contracts, vouchers, or public insurance, can depend on facilities meeting regulatory requirements and quality standards. This strengthens provider payoffs to and incentives for complying with regulatory standards, while channelling patients to free or affordable but higher-quality providers. Again, evaluations should assess the broader effects of interventions, especially on reducing health inequalities.

We have highlighted the paucity of evidence. First, there is no clear evidence on towns and peri-urban areas

where the tendency for providers to crowd in around specialist and training centres might be more attenuated. Second, there is little evidence from the most marginalised urban groups—those who are unregistered or have no fixed abode. Third, LMICs cover a wide range of economic development, and more research and evidence are required on the different needs and opportunities over these very different contexts; the data presented here are weighted towards the low-middle-income stratum. Fourth, we are only at an early stage of developing an evidence base. Many studies evaluate an intervention versus no intervention, but then make no attempt to assess the opportunity cost, and hence value for money. Fifth, few interventions compare alternative types of interventions. Many studies combine intervention types, but then do not examine the effects and costs of individual components and their possible interactions using factorial designs for primary studies or network meta-analysis for systematic reviews. Sixth, it is crucially important to evaluate the whole causal chain connecting intervention to outcome so as to deepen scientific understanding and to estimate how effects will change by context.^{135–137} Seventh, interventions need to evaluate outcomes that go beyond intended outcomes. Finally, most studies come from low-income and low-middle-income countries, with little research from middle-income countries, although insights from the latter, particularly on the development of primary care teams, are highly relevant to lower-income countries.

We might expect a gradual expansion of resources in LMIC health services. First, we might anticipate growth in LMIC economies, even if the rate of growth fluctuates (eg, real yearly economic growth averaged between 4% and 6% across low-income countries between 2001 and 2018).¹³⁸ Second, resource allocations to health care as a whole are set to improve under the impetus of universal health coverage and countries' existing commitments. Third, primary care and health services are moving centre stage. The 2024 WHO report on primary health care summarises the systematic review literature to show that investing in primary care improves health, reduces costs (by reducing the need for hospital treatment), and boosts health equity.⁷² There is also evidence of an economic payback from health care resulting from a more productive workforce and less poverty from catastrophic out-of-pocket payments. We add our voice to the many who advocate for rebalancing towards primary care and prevention, which countries have signed up to in the spirit of the Astana Declaration.¹³⁹

Conclusion

We have reached an optimistic stage in the development of primary care services in LMIC cities. The evidence base on new meso-level strategies is growing rapidly, with new studies evaluating regulatory policies and market-shaping interventions. However, more research is required, especially examining how interventions play

out across the spectrum of LMIC city contexts, evaluating their full causal pathways and cost effectiveness, and exploring their interactions and unintended effects. The need for more research is especially true for the most vulnerable urban populations. Increased future investment is likely and we have suggested potential interventions, not all of which will be equally successful or provide equivalent payback to investment. Cities are the drivers of economic progress—they are where most people in LMICs now live—and the market for primary care services can be reshaped in cities. It is important to develop and evaluate novel approaches to health policies across LMIC cities so that they can realise their potential to create wealth and health.

Contributors

All coauthors contributed to reviewing and editing drafts and approved the final submission. BD: writing the original draft, figures, method, resources, data curation, and formal analysis. BM: conceptualisation, writing the original draft, and method. EPP: conceptualisation and writing the original draft. GA: conceptualisation and literature search. JD: conceptualisation, formal analysis, method, supervision, and writing the original draft. JS: project administration. KabS: conceptualisation. PJC: project administration and visualisation. RJL: conceptualisation, funding acquisition, investigation, methodology, project administration, supervision, visualisation, writing the original draft, and systematic reviewing. SIW: conceptualisation and method. Y-FC: literature search for the review of systematic reviews and advised on the scoping review. ZAB: writing the original draft and method. BD, JD, KScA, RJL, and SIW directly accessed and verified the underlying data reported in this manuscript.

Declaration of interests

RJL, FG, JS, and SIW received funding from National Institute for Health and Care Research (NIHR) Research and Innovation for Global Health Transformation (NIHR 200132) using UK Aid from the UK Government to support global health research. RJL, JS, and SIW received funding from the NIHR Global Health Research Unit on Improving Health in Slums. RJL, EPP, KScA, and CM received funding from the NIHR Global Health Research Unit on Social and Environmental Determinants of Health. RJL, JS, and SIW received funding from NIHR Midlands Patient Safety Research Collaboration. RJL and PJC received funding from NIHR Applied Research Collaboration West Midlands. SIW also received funding from Medical Research Council grant (MR/V038591/1). EPP received support from the Bill & Melinda Gates Foundation to attend symposiums and meetings. CM has a Research Productivity Grant from the National Council for Scientific and Technological Development of Brazil and a Distinguished Scientist Grant from Carlos Chagas Research Foundation—State of Rio de Janeiro. The views expressed in this publication are those of the authors and not necessarily those of the NIHR or the UK Department of Health and Social Care.

References

- 1 Institute of Medicine Committee on Quality of Health Care in America. Crossing the quality chasm: a new health system for the 21st century. National Academies Press, 2001.
- 2 Herrera CA, Lewin S, Paulsen E, et al. Governance arrangements for health systems in low-income countries: an overview of systematic reviews. *Cochrane Database Syst Rev* 2017; 9: CD011085.
- 3 Pantoja T, Opiyo N, Lewin S, et al. Implementation strategies for health systems in low-income countries: an overview of systematic reviews. *Cochrane Database Syst Rev* 2017; 9: CD011086.
- 4 Ciapponi A, Lewin S, Herrera CA, et al. Delivery arrangements for health systems in low-income countries: an overview of systematic reviews. *Cochrane Database Syst Rev* 2017; 9: CD011083.
- 5 Wiysonge CS, Paulsen E, Lewin S, et al. Financial arrangements for health systems in low-income countries: an overview of systematic reviews. *Cochrane Database Syst Rev* 2017; 9: CD011084.
- 6 Rowe AK, Rowe SY, Peters DH, Holloway KA, Chalker J, Ross-Degnan D. Effectiveness of strategies to improve health-care provider practices in low-income and middle-income countries: a systematic review. *Lancet Glob Health* 2018; 6: e1163–75.
- 7 Bitton A, Fifield J, Ratcliffe H, et al. Primary healthcare system performance in low-income and middle-income countries: a scoping review of the evidence from 2010 to 2017. *BMJ Glob Health* 2019; 4 (suppl 8): e001551.
- 8 Angell B, Dodd R, Palagyi A, et al. Primary health care financing interventions: a systematic review and stakeholder-driven research agenda for the Asia-Pacific region. *BMJ Glob Health* 2019; 4 (suppl 8): e001481.
- 9 Dodd R, Palagyi A, Jan S, et al. Organisation of primary health care systems in low- and middle-income countries: review of evidence on what works and why in the Asia-Pacific region. *BMJ Glob Health* 2019; 4 (suppl 8): e001487.
- 10 Bright T, Felix L, Kuper H, Polack S. Systematic review of strategies to increase access to health services among children over five in low- and middle-income countries. *Trop Med Int Health* 2018; 23: 476–507.
- 11 Diaconu K, Falconer J, Verbel A, Fretheim A, Witter S. Paying for performance to improve the delivery of health interventions in low- and middle-income countries. *Cochrane Database Syst Rev* 2021; 5: CD007899.
- 12 Marcolino MS, Oliveira JAQ, D'Agostino M, Ribeiro AL, Alkmim MBM, Novillo-Ortiz D. The impact of mHealth interventions: systematic review of systematic reviews. *JMIR Mhealth Uhealth* 2018; 6: e23.
- 13 Joseph L, Lavis A, Greenfield S, et al. Systematic review on the use of patient-held health records in low-income and middle-income countries. *BMJ Open* 2021; 11: e046965.
- 14 Mash R, Christian C, Chigwanda RV. Alternative mechanisms for delivery of medication in South Africa: a scoping review. *S Afr Fam Pract* 2021; 63: e1–8.
- 15 Watson SI, Sahota H, Taylor CA, Chen Y-F, Lilford RJ. Cost-effectiveness of health care service delivery interventions in low and middle income countries: a systematic review. *Glob Health Res Policy* 2018; 3: 17.
- 16 Papanicolas I, Rajan D, Karanikolos M, Soucat A, Figueras J. Health system performance assessment: a framework for policy analysis. European Observatory on Health Systems and Policies, 2022.
- 17 Lilford RJ, Chilton PJ, Hemming K, Girling AJ, Taylor CA, Barach P. Evaluating policy and service interventions: framework to guide selection and interpretation of study end points. *BMJ* 2010; 341: c4413-c.
- 18 Alonge O, Gupta S, Engineer C, Salehi AS, Peters DH. Assessing the pro-poor effect of different contracting schemes for health services on health facilities in rural Afghanistan. *Health Policy Plan* 2015; 30: 1229–42.
- 19 Elsey H, Agyepong I, Huque R, et al. Rethinking health systems in the context of urbanisation: challenges from four rapidly urbanising low-income and middle-income countries. *BMJ Glob Health* 2019; 4: e001501.
- 20 WHO. What are the overall principles of HBP design? 2024. <https://www.who.int/teams/health-systems-governance-and-financing/economic-analysis/health-technology-assessment-and-benefit-package-design/resource-guide-for-the-use-of-hba-and-hbp-design-processes/what-are-the-overall-principles-of-hbp-design> (accessed Jan 30, 2024).
- 21 Hanson K, Brikci N, Erlangga D, et al. The Lancet Global Health Commission on financing primary health care: putting people at the centre. *Lancet Glob Health* 2022; 10: e715–72.
- 22 Daniels B, Das J, Gatti R, Yi Chang A. Caseloads and competence in sub-Saharan Africa. Policy research working paper #11043. 2025. <https://documents1.worldbank.org/curated/en/09983650132518461/pdf/1DU-673dede9-ed4f-4766-8adf-c804e1275d38.pdf> (accessed March 18, 2025).
- 23 Sturt J, Griffiths F, Ajisola M, et al. Safety and upscaling of remote consulting for long-term conditions in primary health care in Nigeria and Tanzania (REaCH trials): stepped-wedge trials of training, mobile data allowance, and implementation. *Lancet Glob Health* 2023; 11: e1753–64.

- 24 Makene FS, Ngilangwa R, Santos C, et al. Patients' pathways to cancer care in Tanzania: documenting and addressing social inequalities in reaching a cancer diagnosis. *BMC Health Serv Res* 2022; **22**: 189.
- 25 Arsenault C, Rowe SY, Ross-Degnan D, et al. How does the effectiveness of strategies to improve healthcare provider practices in low-income and middle-income countries change after implementation? Secondary analysis of a systematic review. *BMJ Qual Saf* 2022; **31**: 123–33.
- 26 Crispino KT, Rocha ICN. Enhancing healthcare professional practice in the Philippines toward ASEAN integration through the continuing professional development law. *ASEAN Journal of Community Engagement* 2021; **5**: 376–95.
- 27 Mash RJ, Schouw D. Empowerment and Training (GREAT) for diabetes in South Africa: convergent mixed methods. *BMJ Open* 2024; **14**: e085171.
- 28 Liang LB, Li X, Liu XP, et al. Evaluation of the Star Family Doctors Training Program: an observational cohort study of a novel continuing medical education program for general practitioners within a compact medical consortium: a quantitative analysis. *BMC Med Educ* 2023; **23**: 250.
- 29 Ellard DR, Chimwaza W, Davies D, et al. Can training in advanced clinical skills in obstetrics, neonatal care and leadership, of non-physician clinicians in Malawi impact on clinical services improvements (the ETATMBA project): a process evaluation. *BMJ Open* 2014; **4**: e005751.
- 30 Goudge J, Babalola O, Malatji H, Levin J, Thorogood M, Griffiths F. The effect of a roving nurse mentor on household coverage and quality of care provided by community health worker teams in South Africa: a longitudinal study with a before, after and 6 months post design. *BMC Health Serv Res* 2023; **23**: 186.
- 31 Tseng Y-h, Griffiths F, de Kadt J, et al. Integrating community health workers into the formal health system to improve performance: a qualitative study on the role of on-site supervision in the South African programme. *BMJ Open* 2019; **9**: e022186-e.
- 32 Das J, Chowdhury A, Hussam R, Banerjee AV. The impact of training informal health care providers in India: a randomized controlled trial. *Science* 2016; **354**: aaf7384.
- 33 Gadsden T, Mabunda SA, Palagyi A, et al. Performance-based incentives and community health workers' outputs, a systematic review. *Bull World Health Organ* 2021; **99**: 805–18.
- 34 Yousafzai AK, Rasheed MA, Rizvi A, Armstrong R, Bhutta ZA. Effect of integrated responsive stimulation and nutrition interventions in the Lady Health Worker programme in Pakistan on child development, growth, and health outcomes: a cluster-randomised factorial effectiveness trial. *Lancet* 2014; **384**: 1282–93.
- 35 Bhutta ZA, Soofi S, Cousens S, et al. Improvement of perinatal and newborn care in rural Pakistan through community-based strategies: a cluster-randomised effectiveness trial. *Lancet* 2011; **377**: 403–12.
- 36 Eijkenaar F, Emmert M, Scheppach M, Schöffski O. Effects of pay for performance in health care: a systematic review of systematic reviews. *Health Policy* 2013; **110**: 115–30.
- 37 Diaconu K, Witter S, Binyaruka P, et al. Appraising pay-for-performance in healthcare in low- and middle-income countries through systematic reviews: reflections from two teams. *Cochrane Database Syst Rev* 2022; **5**: ED000157.
- 38 de Walque D, Kandpal E. Reviewing the evidence on health financing for effective coverage: do financial incentives work? *BMJ Glob Health* 2022; **7**: e009932.
- 39 Basinga P, Gertler PJ, Binagwaho A, Soucat AL, Sturdy J, Vermeersch CM. Effect on maternal and child health services in Rwanda of payment to primary health-care providers for performance: an impact evaluation. *Lancet* 2011; **377**: 1421–28.
- 40 Kovacs R, Maia Barreto JO, da Silva EN, et al. Socioeconomic inequalities in the quality of primary care under Brazil's national pay-for-performance programme: a longitudinal study of family health teams. *Lancet Glob Health* 2021; **9**: e331–39.
- 41 Kristensen SR, Meacock R, Turner AJ, et al. Long-term effect of hospital pay for performance on mortality in England. *N Engl J Med* 2014; **371**: 540–48.
- 42 Watson SI, Bakibinga P, Kabaria C, et al. Primary care doctor and nurse consultations among people who live in slums: a retrospective, cross-sectional survey in four countries. *BMJ Open* 2022; **12**: e054142-e.
- 43 Schiavetti B, Wynendaele E, De Spiegeleer B, et al. The quality of medicines used in children and supplied by private pharmaceutical wholesalers in Kinshasa, Democratic Republic of Congo: a prospective survey. *Am J Trop Med Hyg* 2018; **98**: 894–903.
- 44 Lamba G, Shroff ZC, Babar Z-U-D, Ghaffar A. Drug shops for stronger health systems: learning from initiatives in six LMICs. *J Pharm Policy Pract* 2021; **14** (suppl 1): 94.
- 45 Sanya RE, Johnston ES, Kibe P, et al. Effectiveness of self-financing patient-led support groups in the management of hypertension and diabetes in low- and middle-income countries: systematic review. *Trop Med Int Health* 2023; **28**: 80–89.
- 46 Vogler S, Fischer S. How to address medicines shortages: findings from a cross-sectional study of 24 countries. *Health Policy* 2020; **124**: 1287–96.
- 47 Diaconu K, Chen YF, Cummins C, Jimenez Moyao G, Manaseki-Holland S, Lilford R. Methods for medical device and equipment procurement and prioritization within low- and middle-income countries: findings of a systematic literature review. *Global Health* 2017; **13**: 59.
- 48 Makwero MK, Majo T, Devarsetty P, et al. Characterising the performance measurement and management system in the primary health care systems of Malawi. *Afr J Prim Health Care Fam Med* 2024; **16**: e1–11.
- 49 Basu S, Andrews J, Kishore S, Panjabi R, Stuckler D. Comparative performance of private and public healthcare systems in low- and middle-income countries: a systematic review. *PLoS Med* 2012; **9**: e1001244.
- 50 Humphries C, Jaganathan S, Panniyammakal J, et al. Investigating discharge communication for chronic disease patients in three hospitals in India. *PLoS One* 2020; **15**: e0230438-e.
- 51 Haggerty JL, Reid RJ, Freeman GK, Starfield BH, Adair CE, McKendry R. Continuity of care: a multidisciplinary review. *BMJ* 2003; **327**: 1219–21.
- 52 Schwarz D, Hirschhorn LR, Kim J-H, Ratcliffe HL, Bitton A. Continuity in primary care: a critical but neglected component for achieving high-quality universal health coverage. *BMJ Glob Health* 2019; **4**: e001435.
- 53 Boule A, Heekes A, Tiffin N, et al. Data centre profile: the provincial health data centre of the Western Cape province, South Africa. *Int J Popul Data Sci* 2019; **4**: 1143.
- 54 Fennelly O, Cunningham C, Grogan L, et al. Successfully implementing a national electronic health record: a rapid umbrella review. *Int J Med Inform* 2020; **144**: 104281.
- 55 Demena BA, Artavia-Mora L, Ouedraogo D, Thiombiano BA, Wagner N. A systematic review of mobile phone interventions (SMS/IVR/calls) to improve adherence and retention to antiretroviral treatment in low- and middle-income countries. *AIDS Patient Care STDs* 2020; **34**: 59–71.
- 56 WHO. WHO guideline recommendations on digital interventions for health system strengthening. 2019. <https://www.who.int/publications/i/item/9789241550505> (accessed March 17, 2025).
- 57 WHO, UNICEF. Operational framework for primary health care: transforming vision into action. 2020. <https://www.who.int/publications/i/item/9789240017832> (accessed March 17, 2025).
- 58 Fawdry R. Paperless records are not in the best interest of every patient. *BMJ* 2013; **346**: f2064.
- 59 Joseph L, Lavis A, Greenfield S, et al. A systematic review of home-based records in maternal and child health for improving informational continuity, health outcomes, and perceived usefulness in low and middle-income countries. *PLoS One* 2022; **17**: e0267192.
- 60 Harris B, Ajisola M, Alam RM, et al. Mobile consulting as an option for delivering healthcare services in low-resource settings in low- and middle-income countries: a mixed-methods study. *Digit Health* 2021; **7**: 20552076211033425.
- 61 Griffiths F, Watkins JA, Huxley C, et al. Mobile consulting (mConsulting) and its potential for providing access to quality healthcare for populations living in low-resource settings of low- and middle-income countries. *Digit Health* 2020; **6**: 2055207620919594.
- 62 Phiri M, Munoriyarwa A. Health chatbots in Africa: scoping review. *J Med Internet Res* 2023; **25**: e35573.

- 63 Xiong S, Palileo-Villanueva LM, Li H, et al. Use of e-health programmes to deliver urban primary health-care services for noncommunicable diseases in middle-income countries. 2021. <https://apo.who.int/publications/i/item/super-use-of-e-health-programmes-to-deliver-urban-primary-health-care-services-for-noncommunicable-diseases-in-middle-income-countries> (accessed March 17, 2025).
- 64 WHO. Transformation of primary health care in Kazakhstan: moving towards a multidisciplinary model. 2023. <https://www.who.int/europe/publications/i/item/WHO-EURO-2023-8269-48041-71196> (accessed March 17, 2025).
- 65 Perez F, Ba H, Dastagire SG, Altmann M. The role of community health workers in improving child health programmes in Mali. *BMC Int Health Hum Rights* 2009; 9: 28.
- 66 Hammack AY, Bickham JN, Gilliard I 3rd, Robinson WT. A community health worker approach for ending the HIV epidemic. *Am J Prev Med* 2021; 61 (suppl 1): S26–31.
- 67 Bastos LSL, Aguilar S, Rache B, et al. Primary healthcare protects vulnerable populations from inequity in COVID-19 vaccination: an ecological analysis of nationwide data from Brazil. *Lancet Reg Health Am* 2022; 14: 100335.
- 68 Wroe EB, Nhlema B, Dunbar EL, et al. A household-based community health worker programme for non-communicable disease, malnutrition, tuberculosis, HIV and maternal health: a stepped-wedge cluster randomised controlled trial in Neno District, Malawi. *BMJ Global Health* 2021; 6: e006535-e.
- 69 WHO. Disease Commodity Packages. 2024. <https://www.who.int/emergencies/disease-commodity-packages> (accessed Jan 25, 2024).
- 70 Mohamoud G, Mash R. The quality of primary care performance in private sector facilities in Nairobi, Kenya: a cross-sectional descriptive survey. *BMC Prim Care* 2022; 23: 120.
- 71 Bresick G, von Pressentin KB, Mash R. Evaluating the performance of South African primary care: a cross-sectional descriptive survey. *S Afr Fam Pract* 2019; 61: 109–16.
- 72 WHO. Implementing the primary health care approach: a primer. 2024. <https://www.who.int/publications/i/item/9789240090583> (accessed March 17, 2025).
- 73 Shi L, Lee DC, Liang H, et al. Community health centers and primary care access and quality for chronically-ill patients—a case-comparison study of urban Guangdong Province, China. *Int J Equity Health* 2015; 14: 90.
- 74 Coovadia H, Jewkes R, Barron P, Sanders D, McIntyre D. The health and health system of South Africa: historical roots of current public health challenges. *Lancet* 2009; 374: 817–34.
- 75 Li X, Krumholz HM, Yip W, et al. Quality of primary health care in China: challenges and recommendations. *Lancet* 2020; 395: 1802–12.
- 76 King JJC, Powell-Jackson T, Makungu C, et al. Effect of a multifaceted intervention to improve clinical quality of care through stepwise certification (SafeCare) in health-care facilities in Tanzania: a cluster-randomised controlled trial. *Lancet Glob Health* 2021; 9: e1262–72.
- 77 Dunsch FA, Evans DK, Eze-Ajoku E, Macis M. Management, supervision, and healthcare: a field experiment. *J Econ Manage Strategy* 2023; 32: 583–606.
- 78 Powell-Jackson T, King JJC, Makungu C, Quaife M, Goodman C. Management practices and quality of care: evidence from the private health care sector in Tanzania. *Econ J* 2023; 134: 436–56.
- 79 Stacey N, Mirelman A, Kreif N, Suhrcke M, Hofman K, Edoaka I. Facility standards and the quality of public sector primary care: evidence from South Africa's "Ideal Clinics" program. *Health Econ* 2021; 30: 1543–58.
- 80 Rowe SY, Ross-Degnan D, Peters DH, Holloway KA, Rowe AK. The effectiveness of supervision strategies to improve health care provider practices in low- and middle-income countries: secondary analysis of a systematic review. *Hum Resour Health* 2022; 20: 1.
- 81 Gilson L, Daire J. Leadership and governance within the South African health system. *S Afr Health Rev* 2011; 2011: 69–80.
- 82 Nkomazana O, Mash R, Wojczewski S, Kutalek R, Phaladze N. How to create more supportive supervision for primary healthcare: lessons from Ngamiland district of Botswana: co-operative inquiry group. *Glob Health Action* 2016; 9: 31263.
- 83 Schriver M, Cubaka VK, Nyirazinyoye L, Itangishaka S, Kallestrup P. The relationship between primary healthcare providers and their external supervisors in Rwanda. *Afr J Prim Health Care Fam Med* 2017; 9: e1–11.
- 84 Bailey C, Blake C, Schriver M, Cubaka VK, Thomas T, Martin Hilber A. A systematic review of supportive supervision as a strategy to improve primary healthcare services in sub-Saharan Africa. *Int J Gynaecol Obstet* 2016; 132: 117–25.
- 85 Mash R. The contribution of family physicians to African health systems. *Afr J Prim Health Care Fam Med* 2022; 14: e1–9.
- 86 Mash RJ, Govender S, Isaacs AA, De Sa A, Schlemmer A. An assessment of organisational values, culture and performance in Cape Town's primary healthcare services. *S Afr Fam Pract* 2013; 55: 459–66.
- 87 Mash R, Blitz J, Malan Z, Von Pressentin K. Leadership and governance: learning outcomes and competencies required of the family physician in the district health system. *S Afr Fam Pract* 2016; 58: 232–35.
- 88 Lilford R, Omigbodun A, Umar UA, et al. Reducing delays in cancer care in sub-Saharan Africa. <https://www.dev.fundingawards.nih.ac.uk/award/NIHR158242> (accessed March 17, 2025).
- 89 Lagarde M, Palmer N. The impact of user fees on access to health services in low- and middle-income countries. *Cochrane Database Syst Rev* 2011; 2011: CD009094.
- 90 Powell-Jackson T, Hanson K, Whitty CJM, Ansah EK. Who benefits from free healthcare? Evidence from a randomized experiment in Ghana. *J Dev Econ* 2014; 107: 305–19.
- 91 Watson SI, Wroe EB, Dunbar EL, et al. The impact of user fees on health services utilization and infectious disease diagnoses in Neno District, Malawi: a longitudinal, quasi-experimental study. *BMC Health Serv Res* 2016; 16: 595.
- 92 Lochoro P, Bataringaya J, Tashobya CK, Kyabaggu JH. Public-private partnership in health: working together to improve health sector performance in Uganda. In: Tashobya CK, Ssengooba F, Cruz VO, eds. Health systems reforms in Uganda: processes and outputs. London School of Hygiene & Tropical Medicine, 2006: 83–96.
- 93 Bronsoler A, Gruber J, Seira E. The impact of a private supplement to public health care: the Mexico Diabetes Experiment. 2021. <http://www.nber.org/papers/w28913> (accessed March 17, 2025).
- 94 Nachtnebel M, O'Mahony A, Pillai N, Hort K. Effectively engaging the private sector through vouchers and contracting—a case for analysing health governance and context. *Soc Sci Med* 2015; 145: 193–200.
- 95 Albis MLF, Bhadra SK, Chin B. Impact evaluation of contracting primary health care services in urban Bangladesh. *BMC Health Serv Res* 2019; 19: 854.
- 96 Nandraj S. Reconfiguring urban primary healthcare operations, utilisation and financing of Mohalla Clinics of Delhi. *Econ Politic Week* 2021; 25: 112–19.
- 97 Ezech A, Oyebo O, Satterthwaite D, et al. The history, geography, and sociology of slums and the health problems of people who live in slums. *Lancet* 2017; 389: 547–58.
- 98 Bakibinga P, Ettarh R, Ziraba AK, et al. The effect of enhanced public-private partnerships on maternal, newborn and child health Services and outcomes in Nairobi-Kenya: the PAMANECH quasi-experimental research protocol. *BMJ Open* 2014; 4: e006608.
- 99 Abel-Smith B, Dua A. Community-financing in developing countries: the potential for the health sector. *Health Policy Plan* 1988; 3: 95–108.
- 100 Garner P. The Bamako Initiative. *BMJ* 1989; 299: 277–78.
- 101 McPake B, Hanson K, Mills A. Community financing of health care in Africa: an evaluation of the Bamako initiative. *Soc Sci Med* 1993; 36: 1383–95.
- 102 Fouakeng F, Njoumeme Z, Mbacham WF, Angwafo Iii F. Community-based mutual aid for improving financial access to healthcare: case study of 150 community-based organizations in Cameroon. *Int J Community Med Public Health* 2022; 9: 1257–62.
- 103 Jain A, Swetha S, Johar Z, Raghavan R. Acceptability of, and willingness to pay for, community health insurance in rural India. *J Epidemiol Glob Health* 2014; 4: 159–67.

- 104 Victor BWM, Alonga BM, Simbananiye L, Mwilarhe PA, Sindayigaya I. Organisational health and resilience of community health insurance schemes in Bukavu eastern DRC. *Open J Soc Sci* 2023; **11**: 378–98.
- 105 Dake FAA. Examining equity in health insurance coverage: an analysis of Ghana's National Health Insurance Scheme. *Int J Equity Health* 2018; **17**: 85.
- 106 Jakab M, Krishnan C. Community involvement in health care financing: a survey of the literature on the impacts, strengths and weaknesses. World Bank, 2001.
- 107 Putri NK, Laksono AD, Rohmah N. Predictors of national health insurance membership among the poor with different education levels in Indonesia. *BMC Public Health* 2023; **23**: 373.
- 108 Shrestha D, Napit IB, Ansari S, et al. Evaluation of a self-help intervention to promote the health and wellbeing of marginalised people including those living with leprosy in Nepal: a prospective, observational, cluster-based, cohort study with controls. *BMC Public Health* 2021; **21**: 873.
- 109 Richterman A, Millien C, Bair EF, et al. The effects of cash transfers on adult and child mortality in low- and middle-income countries. *Nature* 2023; **618**: 575–82.
- 110 Pega F, Pabayo R, Benny C, Lee E-Y, Lhachimi SK, Liu SY. Unconditional cash transfers for reducing poverty and vulnerabilities: effect on use of health services and health outcomes in low- and middle-income countries. *Cochrane Database Syst Rev* 2022; **3**: CD011135.
- 111 Onuegbu C, Larweh M, Harlock J, Griffiths F. Systematic review of lay consultation in symptoms and illness experiences in informal urban settlements of low-income and middle-income countries. *BMJ Open* 2021; **11**: e050766.
- 112 Prost A, Colbourn T, Seward N, et al. Women's groups practising participatory learning and action to improve maternal and newborn health in low-resource settings: a systematic review and meta-analysis. *Lancet* 2013; **381**: 1736–46.
- 113 Louart S, Bonnet E, Ridde V. Is patient navigation a solution to the problem of "leaving no one behind"? A scoping review of evidence from low-income countries. *Health Policy Plan* 2021; **36**: 101–16.
- 114 WHO. WHO global strategy on people-centred and integrated health services. Interim report. 2015. <https://iris.who.int/handle/10665/155002> (accessed March 17, 2025).
- 115 Montagu D, Goodman C. Prohibit, constrain, encourage, or purchase: how should we engage with the private health-care sector? *Lancet* 2016; **388**: 613–21.
- 116 The Health Foundation. Evidence scan: competition in healthcare. 2011. <https://www.health.org.uk/sites/default/files/CompetitionInHealthcare.pdf>. (accessed March 17, 2025).
- 117 Bedoya G, Das J, Dolinger A. Randomized regulation: the impact of minimum quality standards on health markets. Policy Research Working Paper; 10386. 2023. <http://hdl.handle.net/10986/39630> (accessed March 17, 2025).
- 118 McPake B, Hanson K. Managing the public-private mix to achieve universal health coverage. *Lancet* 2016; **388**: 622–30.
- 119 Mulcahy P, Mahal A, McPake B, Kane S, Ghosh PK, Lee JT. Is there an association between public spending on health and choice of healthcare providers across socioeconomic groups in India?—evidence from a national sample. *Soc Sci Med* 2021; **285**: 114149.
- 120 Lim KS. Does the market correct political distortions? Evidence from hospital construction in Malaysia. Harvard University, 2023.
- 121 Flodgren G, Pomey MP, Taber SA, Eccles MP. Effectiveness of external inspection of compliance with standards in improving healthcare organisation behaviour, healthcare professional behaviour or patient outcomes. *Cochrane Database Syst Rev* 2011; **9**: CD008992.
- 122 Flodgren G, Gonçalves-Bradley DC, Pomey MP. External inspection of compliance with standards for improved healthcare outcomes. *Cochrane Database Syst Rev* 2016; **12**: CD008992.
- 123 Stenson B, Syhakhang L, Lundborg CS, Eriksson B, Tomson G. Private pharmacy practice and regulation. A randomized trial in Lao P.D.R. *Int J Technol Assess Health Care* 2001; **17**: 579–89.
- 124 Chalker J, Ratanawijitrasin S, Chuc NT, Petzold M, Tomson G. Effectiveness of a multi-component intervention on dispensing practices at private pharmacies in Vietnam and Thailand—a randomized controlled trial. *Soc Sci Med* 2005; **60**: 131–41.
- 125 Jacobs TG, Robertson J, van den Ham HA, Iwamoto K, Bak Pedersen H, Mantel-Teeuwisse AK. Assessing the impact of law enforcement to reduce over-the-counter (OTC) sales of antibiotics in low- and middle-income countries: a systematic literature review. *BMC Health Serv Res* 2019; **19**: 536.
- 126 Johnson MC, Schellekens O, Stewart J, van Ostenberg P, de Wit TR, Spieker N. SafeCare: an innovative approach for improving quality through standards, benchmarking, and improvement in low- and middle-income countries. *Jt Comm J Qual Patient Saf* 2016; **42**: 350–71.
- 127 Quinn L, Bird P, Remsing S, Reeves K, Lilford R. Unintended consequences of the 18-week referral to treatment standard in NHS England: a threshold analysis. *BMJ Qual Saf* 2023; **32**: 712–20.
- 128 Damrich S, Kealey T, Ricketts M. Crowding in and crowding out within a contribution good model of research. *Res Policy* 2022; **51**: 104400.
- 129 Bryan G, Glaeser E, Tsivanidis N. Cities in the developing world. *Annu Rev Econ* 2020; **12**: 273–97.
- 130 Das J, Do Q-T. The prices in the crises: what we are learning from twenty years of health insurance in low- and middle-income countries. *J Econ Perspect* 2023; **37**: 123–52.
- 131 Schneider H, Masilela T, Mndelebe J, et al. Special series on the district health system. *S Afr Med J* 2023; **113**: 14.
- 132 Marcil L, Afsana K, Perry HB. First steps in initiating an effective maternal, neonatal, and child health program in urban slums: the BRAC Manoshi project's experience with community engagement, social mapping, and census taking in Bangladesh. *J Urban Health* 2016; **93**: 6–18.
- 133 More NS, Das S, Bapat U, et al. Community resource centres to improve the health of women and children in informal settlements in Mumbai: a cluster-randomised, controlled trial. *Lancet Glob Health* 2017; **5**: e335–49.
- 134 Habib MA, Soofi SB, Hussain Z, et al. A holistic strategy of mother and child health care to improve the coverage of routine and polio immunization in Pakistan: results from a demonstration project. *Vaccines* 2024; **12**: 89.
- 135 Skivington K, Matthews L, Simpson SA, et al. A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance. *BMJ* 2021: n2061-n.
- 136 Sutton M, Garfield-Birkbeck S, Martin G, et al. Economic analysis of service and delivery interventions in health care. *Health Serv Deliv Res* 2018; **6**: 1–16.
- 137 Watson SI, Lilford RJ. Integrating multiple sources of evidence: a Bayesian perspective. *Health Serv Deliv Res* 2018; **4**: 1–14.
- 138 Steinbach R. Growth in low-income countries: evolution, prospects, and policies. The World Bank, 2019.
- 139 WHO. Declaration of Astana. 2018. <https://www.who.int/publications/i/item/WHO-HIS-SDS-2018.61> (accessed March 17, 2025).

Copyright © 2025 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.